

**52 исследования
эффективности состава
ОптиФайбер**

OptiFibre®

52 исследования эффективности состава ОптиФайбер

1. Takeno F, Yamada H, Sekiya K, Fujitani B, Ohtsu K. Effect of partially decomposed guar gum on high-cholesterol-fed rats and non-dietary fiber-fed rats. *JJpnSocNutrFoodSci*. 1990;43(6):421-425.
2. Ide T, Mariuchi H, Nihimoto K. Hypolipidemic effects of guar gum and its enzyme hydrolysate in rats fed highly saturated fat diets. *Ann Nutr Metab*. 1991;35:34-44.
3. Lampe JW, Effertz ME, Larson JL, Slavin JL. Gastrointestinal effects of modified guar gum and soy polysaccharide as part of an enteral formula diet. *JPEN*. 1992;16(6):538-544.
4. Yamatoya K, Sekiya K, Yamada H, Ichikawa T. Effects of partially hydrolysed guar gum on postprandial plasma glucose and lipid levels in humans. *JJpnSocNutrFoodSci*. 1993;46(3):199-203.
5. Takahashi H, Yang SI, Hayashi C, Kim M, Yamanaka J, Yamamoto T. Effect of partially hydrolyzed guar gum on fecal output in human volunteers. *NutrRes*. 1993;13:649-657.
6. Meier R, Beglinger C, Schneider H, Rowdner A, Gyr K. Effect of liquid diet with and without soluble fiber supplementation on intestinal transit and cholecystokinin release in volunteers. *JPEN*. 1993;17:231-235.
7. Takahashi H, Yang SI, Kim M, Yamamoto T. Protein and energy utilization of growing rats fed on the diets containing intact or partially hydrolyzed guar gum. *CompBiochemPhysiol*. 1994;107A(1):255-260.
8. Van de Ven ML, Westerterp-Plantenga MS, Wouters L, Saris WH. Effects of liquid preloads with different fructose/fiber concentrations on subsequent food intake and ratings of hunger in women. *Appetite*. 1994;23:139-146.
9. Takahashi H, Wako N, Okubo T, Ishihara N, Yamanaka J, Yamamoto T. Influence of partially hydrolyzed guar gum on constipation in women. *J Nutr Sci Vitaminol*. 1994;40:251-259.
10. Takahashi H, Yang SI, Ueda Y, Kim M, Yamamoto T. Influence of intact and partially hydrolysed guar gum on iron utilization in rats fed on iron-deficient diets. *CompBiochem Physiol*. 1994;109A(1):75-82.



52 исследования эффективности состава ОптиФайбер

11. Takahashi H, Yang SI, Fujiki M, Kim M, Yamamoto T, Greenberg NA. Toxicity studies of partially hydrolyzed guar gum. *J Am College of Toxicol.* 1994;13:273-278.
12. Okubo T, Ishihara N, Takahashi H, Fujisawa T, Kim M, Yamamoto T, Mitsuoka T. Effects of partially hydrolyzed guar gum intake on human intestinal microflora and its metabolism. *Biosci Biotechnol Biochem.* 1994;58(8):1364-1369.
13. Homann HH, Kemen M, Fuesslich C, Senkai M, Zumtobel V. Reduction in diarrhea incidence by soluble fiber in patients receiving total or supplemental enteral nutrition. *JPEN.* 1994;18(6):486-490.
14. Takahashi H, Akachi S, Ueda Y, Akachi S, Kim M, Hirano K, Yamamoto T. Effect of liquid diets with or without partially hydrolyzed guar gum on intestinal function of rats. *Nutr Res.* 1995;15(4):527-536.
15. Golay A, Schneider H, Bloise D, Vadas L, Assal JPh. The effect of a liquid supplement containing guar gum and fructose on glucose tolerance in non-insulin-dependent diabetic patients. *Nutr Metab Cardiovasc Dis.* 1995;5:141-148.
16. Yamatoya K, Kuwano K, Suzuki J, Mitamura T, Sekiya K. Effect of hydrolyzed guar gum on frequency and feeling of defecation in humans. *Oyo Toshitsu Kagaku.* 1995;42(3):251-257.
17. Furuse M, Mabayo RT. Effects of partially hydrolysed guar gum on feeding behaviour and crop emptying rate in chicks. *British Poultry Science.* 1996;37:223-227.
18. Hara H, Nagata M, Ohta A, Kasai T. Increases in calcium absorption with ingestion of soluble dietary fibre, guar gum hydrolysate, depend on the caecum in partially nephrectomized and normal rats. *Br J Nutr.* 1996;76:773-784.
19. Weaver GA, Tangel CT, Krause JA, Alpern HD, Jenkins PL, Parfitt MM, Stragand JJ. Dietary guar gum alters colonic microbial fermentation in azoxymethane-treated rats. *J Nutr.* 1996;126:1979-1991.
20. Peters AL, Davidson MB. Addition of hydrolyzed guar to enteral feeding products in type 1 diabetic patients. *Diabetes Care.* 1996;19(8):899-900.



52 исследования эффективности состава ОптиФайбер

21. Yamatoya K, Kuwano K, Suzuki J. Effects of hydrolyzed guar gum on cholesterol and glucose in humans. *Food Hydrocolloids*. 1997;11(2):239-242.
22. Favier ML, Bost PE, Guittard C, Demigne C, Remesy C. The cholesterol-lowering effect of guar gum is not the result of a simple diversion of bile acids toward fecal excretion. *Lipids*. 1997;32(9):953-959.
23. Greenberg NA, Sellman D. Partially hydrolyzed guar gum as a source of fiber. *Cereal Foods World*. 1998;43(9):703-707.
24. Tsuda K, Inden T, Yamanaka K, Ikeda Y. Effect of partially hydrolyzed guar gum on elevation of blood glucose after sugar intake in human volunteers. *J Japan Assoc Diet Fiber Res*. 1998;2(1):15-22.
25. Heini AF, Lara-Castro C, Schneider H, Kirk KA, Considine RV, Weinsier RL. Effect of hydrolyzed guar fiber on fasting and postprandial satiety and satiety hormones: a double-blind, placebo-controlled trial during controlled weight loss. *Int J Obesity*. 1998;22:906-909.
26. Alam NH, Meier R, Rausch T, Meyer-Wyss B, Hildebrand P, Schneider H, Bachmann C, Minder E, Fowler B, Gyr K. Effects of a partially hydrolyzed guar gum on intestinal absorption of carbohydrate, protein and fat: a double-blind controlled study in volunteers. *Clin Nutr*. 1998;17:125-129.
27. Patrick PG, Gohman SM, Marx SC, CeLegge MH, Greenberg NA. Effect of supplements of partially hydrolyzed guar gum on the occurrence of constipation and use of laxative agents. *Research and Professional Briefs*. 1998;98(8):912-914.
28. Yamada K, Tokunaga Y, Ikeda A, Ohkura K, Mamiya S, Kaku S, Sugano M, Tachibana H. Dietary effect of guar gum and its partially hydrolyzed product on the lipid metabolism and immune function of Sprague-Dawley rats. *Biosci Biotechnol Biochem*. 1999;63(12):2163-2167.
29. Hara H, Suzuki T, Kasai T, Aoyama Y, Ohta A. Ingestion of guar gum hydrolysate, a soluble fiber, increases calcium absorption in totally gastrectomized rats. *J Nutr*. 1999;129(1):39-45.



52 исследования эффективности состава ОптиФайбер

30. Hara H, Suzuki T, Kasai T, Aoyama Y, Ohta A. Ingestion of guar-gum hydrolysate partially restores calcium absorption in the large intestine lowered by suppression of gastric acid secretion in rats. *Br J Nutr.* 1999;81:315-321.
31. Valazquez M, Davies C, Marett R, Slavin JL, Feirtag JM. Effect of oligosaccharides and fiber substitutes on short-chain fatty acid production by human faecal microflora. *Anaerobe.* 2000;6:87-92.
32. Watanabe O, Hara H, Aoyama Y, Kasai T. Increased intestinal calcium absorption for the ingestion of a phosphorylated guar gum hydrolysate independent of cecal fermentation in rats. *Biosci Biotechnol Biochem.* 2000;64(3):613-616.
33. Alam NH, Meier R, Schneider H, Sarker SA, Bardhm PK, Mahalanabis D, Fuchs GJ, Gyr N. Partially hydrolyzed guar gum-supplemented oral rehydration solution in the treatment of acute diarrhea in children. *JPediatr Gastroenterol Nutr.* 2000;31:503-507.
34. Watanabe O, Hara H, Kasai T. Effect of a phosphorylated guar gum hydrolysate on increased calcium solubilization and the promotion of calcium absorption in rats. *Biosci Biotechnol Biochem.* 2000;64(1):160-166.
35. Tuohy KM, Kolida S, Lustenberger AM, Gibson GR. The prebiotic effects of biscuits containing partially hydrolysed guar gum and fructo-oligosaccharides – a human volunteer study. *Br J Nutr.* 2001;86:341-348.
36. Spapen H, Diltoer M, Van Maldere C, Opdenaker G, Suys E, Huyghens L. Soluble fiber reduces the incidence of diarrhea in septic patients receiving total enteral nutrition: a prospective, double-blind, randomized and controlled trial. *Clin Nutr.* 2001;20(4):301-305.
37. Nakao M, Ogura Y, Satake S, Ito I, Iguchi A, Takagi K, Nabeshima T. Usefulness of soluble dietary fiber for the treatment of diarrhea during enteral nutrition in elderly patients. *Nutrition.* 2002;18:35-39.



52 исследования эффективности состава ОптиФайбер

38. Parisi GC, Zilli M, Miani MP, Carrara M, Bottona E, Verdianelli G, Battaglia G, Desideri S, Faedo A, Marzolino C, Tonon A, Ermani M, Leandro G. High-fiber diet supplementation in patients with irritable bowel syndrome (IBS): A multicenter, randomized, open trial comparison between wheat bran diet and partially hydrolyzed guar gum (PHGG). *DigDis Sci.* 2002;47(8):1697-1704.
39. Wolf BW, Wlewer TMS, Lai CS, Bolognesi C, Radmard R, Maharry KS, Garleb KA, Hertzler SR, Firkins JL. Effects of a beverage containing an enzymatically induced viscosity dietary fiber, with or without fructose, on the postprandial glycemic response to a high glycemic index food in humans. *Eur J Clin Nutr.* 2003;57:1120-1127.
40. Yamada K, Tokunaga Y, Ikeda A, Ohkura K, Kaku-ohkura S, Mamiya S, Lim BO, Tachibana H. Effect of dietary fiber on the lipid metabolism and immune function of aged Sprague-Dawley rats. *Biosci Biotechnol Biochem.* 2003;67(2):429-433.
41. Slavin JL, Greenberg NA. Partially hydrolyzed guar gum: clinical nutrition uses. *Nutrition.* 2003;19:549-552.
42. Trinidad T, Perez E, Loyola A, Mallillin A, Encabo R, Yokawa T, Aoyama N, Juneja L. Glycemic index of Sunfiber (Cyamopsis tetragonolobus) products in normal and diabetic subjects. *International Journal of Food Science and Technology.* 2004;39:1093-1098.
43. Bosaeus I. Fibre effects of intestinal functions (diarrhoea, constipation and irritable bowel syndrome). *Clinical Nutrition Supplements.* 2004;1:33-38.
44. Kondo S, Xiao J, Takahashi N, Miyaji K, Iwatsuki K, Kokubo S. Suppressing effects of dietary fiber in yogurt on the postprandial serum lipid levels of adult male volunteers. *Biosci Biotechnol Biochem.* 2004;68(5):1135-1138.
45. Bar A. Reducing the glycemic impact of food – a new role for some dietary fibres. *Innovations in Food Technology.* February 2004;34-38.



52 исследования эффективности состава ОптиФайбер

46. Suzuki T, Hara H. Ingestion of guar gum hydrolysate, a soluble and fermentable nondigestible saccharide, improves glucose intolerance and prevents hypertriglyceridemia in rats fed fructose. *J Nutr.* 2004;134:1942-1947.
47. Williams JA, Lai CS, Corwhin H, Ma Y, Maki KC, Garleb KA, Wolf BW. Inclusion of guar gum and alginate into a crispy bar improves postprandial glycemia in humans. *J Nutr.* 2004;134:886-889.
48. Rushdi TA, Pichard C, Khater YH. Control of diarrhea by fiber-enriched diet in ICU patients on enteral nutrition: a prospective randomized controlled trial. *Clin Nutr.* 2004;23:1344-1352.
49. Alam NH, Meier R, Sarker SA, Bardhan PK, Schneider H, Gyr N. Partially hydrolysed guar gum supplemented comminuted chicken diet in persistent diarrhoea: a randomised controlled trial. *Arch Dis Child.* 2005;90(2):195-199.
50. American Heart Association. Online at: www.americanheart.org. Consulted on March 1, 2005.
51. American Diabetes Association. Online at www.diabetes.org . Consulted on March 1, 2005.
52. Dietary Guidelines for Americans 2005. Published jointly by the Department of Health and Human Services (HHS) and the Department of Agriculture (USDA). Online at: <http://www.healthierus.gov/dietaryguidelines/> Consulted on March 1, 2005.

